



US006602672B1

(12) **United States Patent**
Prusiner et al.

(10) **Patent No.: US 6,602,672 B1**
(45) **Date of Patent: Aug. 5, 2003**

(54) **RECOMBINANT CONSTRUCT ENCODING
EPI TOPE TAGGED PRP PROTEIN**

(75) Inventors: **Stanley B. Prusiner**, San Francisco, CA (US); **Glenn C. Telling**, San Francisco, CA (US); **Fred E. Cohen**, San Francisco, CA (US); **Michael R. Scott**, San Francisco, CA (US)

(73) Assignee: **The Regents of the University of California**, Oakland, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 25 days.

(21) Appl. No.: **09/669,516**

(22) Filed: **Sep. 25, 2000**

Related U.S. Application Data

(63) Continuation of application No. 09/031,168, filed on Feb. 26, 1998, now Pat. No. 6,150,583, which is a division of application No. 08/660,626, filed on Jun. 6, 1996, now Pat. No. 5,789,655, which is a continuation-in-part of application No. 08/521,992, filed on Aug. 31, 1995, now Pat. No. 5,908,969, which is a continuation-in-part of application No. 08/509,261, filed on Jul. 31, 1995, now Pat. No. 5,763,740, which is a continuation-in-part of application No. 08/242,188, filed on May 13, 1994, now Pat. No. 5,565,186.

(51) **Int. Cl.**⁷ **G01N 33/53**; C12P 21/06; C07H 21/02; C07K 14/00

(52) **U.S. Cl.** **435/7.1**; 435/69.1; 435/320.1; 435/325; 435/252.3; 536/23.1; 530/350

(58) **Field of Search** 536/23.1; 435/325, 435/252.3, 320.1, 7.1, 69.1; 530/350

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,873,191	A	10/1989	Wagner et al.
5,487,992	A	1/1996	Capecchi et al.
5,545,808	A	8/1996	Hew et al.
5,554,512	A	9/1996	Lyman et al.
5,565,186	A	10/1996	Prusiner et al.
5,789,655	A	8/1998	Prusiner et al.

FOREIGN PATENT DOCUMENTS

WO	WO 94/16092	7/1994
WO	WO 95/20666	8/1995
WO	WO 95/31466	11/1995

OTHER PUBLICATIONS

Basler, K., et al., (1986) "Scrapie and cellular PrP isoforms are encoded by the same chromosomal gene", *Cell* 46:417-428.

Bolton, D.C., et al., (1982) "Identification of a protein that purifies with the scrapie prion", *Science* vol. 218:1309-1311.

Borchelt, D.R., et al., (1990) "Scrapie and cellular prion proteins differ in their kinetics of synthesis and topology in cultured cells." *J. Cell Biol.* 110:743-52.

Bueler, H., et al., (1992) "Normal development and behavior of mice lacking the neuronal cell-surface PrP protein", *Nature* 356.

Gabriel, J.-M., et al., (1992) "Molecular cloning of a candidate chicken prion protein", *Proc. Natl. Acad. Sci. USA* 89:9097-9101.

Hecker, R., et al., "Replication of distinct scrapie prion isolates is region specific in brains of transgenic mice and hamsters", *Genes and Development* 6:1213-1228.

Houdebine, Louis-Marie, (1994), "Production of Pharmaceutical Proteins from Transgenic Animals", *J. of Biotechnology* 34:269-287.

Kascsak, R.J., et al., (1987) "Mouse polyclonal and monoclonal antibody to scrapie-associated fibril proteins", *J. Virol.* 61:3688-93.

Knappik, A., and Pluckthun, A., (1994) "An improved affinity tag based on the FLAG peptide for the detection and purification of recombinant antibody fragments", *Biotechniques* 17:754-61.

Kunz, D., et al., (1992) "The human leukocyte platelet-activating factor receptor", *J. Biol. Chem.* 267:9101-6.

McKinley, P., et al., (1983) "A protease-resistant protein is a structural component of the scrapie prion", *Cell* 36:57-62.

Murray, P.J., et al., (1995) "Epitope tagging of the human endoplasmic reticulum HSP70 protein, BiP, to facilitate Analysis of BiP-substrate interactions", *Anal Biochem.* 229:170-9.

Pan, K.-M., et al., (1993) "Conversion of α -helicies into β -sheets features in the formation of the scrapie prion proteins", *Proc. Natl. Acad. Sci. USA* 90:10962-6.

Prusiner, et al., (1993) "Ablation of the Prion Protein (PrP) Gene in Mice Prevents Scrapie and Facilitates Production of Anti-PrP Antibodies", *Proc. Natl. Acad. Sci. USA* 90:10608-12.

Prusiner, S.B., et al., (1991) "Molecular Biology of prion diseases", *Sciences* 252:1515-1522.

Pursel, et al., (1989) "Genetic Engineering Livestock", *Science* 244:1281-8.

Ruoslahti, E., and Pierschbacher, M.D., (1987) "New perspectives in cell adhesion: RGD and integrins", *Science* 238:491-7.

Salter, et al., (1987) "Transgenic Chickens: Insertion of Retroviral Genes into the Chicken Germ Line", *Virol.* 157:236-40.

(List continued on next page.)

Primary Examiner—Karen Cochrane Carlson
(74) **Attorney, Agent, or Firm**—Karl Bozicevic; Bozicevic, Field & Francis LLP

(57) **ABSTRACT**

DNA constructs are provided of epitope-tagged proteins or protein fragments which are conveniently purified with immunoaffinity chromatography such as epitope-tagged prion proteins (PrP). Transgenic animals expressing an epitope-tagged protein are provided, including transgenic animals expressing epitope-tagged PrP. Methods for distinguishing between the conformational shapes of a protein and a convenient method for isolating a tagged protein by immunoaffinity chromatographic methods are provided.

18 Claims, 6 Drawing Sheets

OTHER PUBLICATIONS

Schmidt, T.G.M., and Skerra, A., (1994) "One-step affinity purification of bacterially produced proteins by means of the "Strep tag" and immobilized recombinant core streptavidin", *J. Chromatography* 676:337-45.

Scott, M.R., et al., (1989) "Chimeric prion protein expression in cultured cells and transgenic mice", *Cell* 59:847-57.

Scott, M., et al., (1992) "Transgenic mice expressing hamster prion protein produce species-specific scrapie infectivity and amyloid plaques", *Protein Sci.* 1:986-997.

Telling, G.C., et al., (1995) "Prion propagation in mice expressing human and chimeric PrP transgenes implicates

the interaction of cellular PrP with another protein", *Cell* 83:79-90.

Telling et al., (1994) "Transmission of Creutzfeldt-Jakob Disease from Humans to Transgenic Mice Expressing Chimeric Human-Mouse Prion Protein", *Proc. Natl. Acad. Sci. USA* 91:9936-40.

Westway. D., et al., (1994) "Degeneration of Skeletal Muscle, Peripheral Nerves, and the Central Nervous system in transgenic mice overexpressing wild-type prion proteins", *Cell* 76:117-29.

FIG. 1

Mo	Met	Ala	Asn	Leu	Gly	Tyr	Trp	Leu	Leu	Ala	Leu	Phe	Val	Thr	Met	Trp	16
Hu						Cys		Met		Val				Ala	Thr		
Mo	Thr	Asp	Val	Gly	Leu	Cys	Lys	Lys	Arg	Pro	Lys	Pro	Gly	Gly	Trp	Asn	32
Hu	Ser		Leu														
Mo	Thr	Gly	Gly	Ser	Arg	Tyr	Pro	Gly	Gln	Gly	Ser	Pro	Gly	Gly	Asn	Arg	48
Hu																	
Mo	Tyr	Pro	Pro	Gln	Gly	Gly	---	Thr	Trp	Gly	Gln	Pro	His	Gly	Gly	Gly	63
Hu							Gly	Gly									
Mo	Trp	Gly	Gln	Pro	His	Gly	Gly	Ser	Trp	Gly	Gln	Pro	His	Gly	Gly	Ser	79
Hu								Gly								Gly	
Mo	Trp	Gly	Gln	Pro	His	Gly	Gly	Gly	Trp	Gly	Gln	Gly	Gly	Gly	Thr	His	95
Hu																	
Mo	Asn	Gln	Trp	Asn	Lys	Pro	Ser	Lys	Pro	Lys	Thr	Asn	Leu	Lys	His	Val	111
Hu	Ser												Met			Met	
Mo	Ala	Gly	Ala	Ala	Ala	Ala	Gly	Ala	Val	Val	Gly	Gly	Leu	Gly	Gly	Tyr	127
Hu																	
Mo	Met	Leu	Gly	Ser	Ala	Met	Ser	Arg	Pro	Met	Ile	His	Phe	Gly	Asn	Asp	143
Hu										Ile					Ser		
Mo	Trp	Glu	Asp	Arg	Tyr	Tyr	Arg	Glu	Asn	Met	Tyr	Arg	Tyr	Pro	Asn	Gln	159
Hu	Tyr										His						
Mo	Val	Tyr	Tyr	Arg	Pro	Val	Asp	Gln	Tyr	Ser	Asn	Gln	Asn	Asn	Phe	Val	175
Hu						Met		Glu									
Mo	His	Asp	Cys	Val	Asn	Ile	Thr	Ile	Lys	Gln	His	Thr	Val	Thr	Thr	Thr	191
Hu																	
Mo	Thr	Lys	Gly	Glu	Asn	Phe	Thr	Glu	Thr	Asp	Val	Lys	Met	Met	Glu	Arg	207
Hu																	
Mo	Val	Val	Glu	Gln	Met	Cys	Val	Thr	Gln	Tyr	Gln	Lys	Glu	Ser	Gln	Ala	223
Hu							Ile				Glu	Arg					
Mo	Tyr	Tyr	Asp	Gly	Arg	Arg	Ser	Ser	Ser	Thr	Val	Leu	Phe	Ser	Ser	Pro	239
Hu			Gln	---	---		Gly			Met							
Mo	Pro	Val	Ile	Leu	Leu	Ile	Ser	Phe	Leu	Ile	Phe	Leu	Ile	Val	Gly		254
Hu																	

(SEQ ID NO: 8)
 (SEQ ID NO: 7)

FIG. 2

```

Mo Met Ala Asn Leu --- --- Gly Tyr Trp Leu Leu Ala Leu Phe Val Thr 14
Bo      Val Lys Ser His Ile      Ser      Ile      Val      Ala

Mo Met Trp Thr Asp Val Gly Leu Cys Lys Lys Arg Pro Lys Pro Gly Gly 30
Bo              Ser

Mo --- Trp Asn Thr Gly Gly Ser Arg Tyr Pro Gly Gln Gly Ser Pro Gly 45
Bo

Mo Gly Asn Arg Tyr Pro Pro Gln Gly Gly --- Thr Trp Gly Gln Pro His 60
Bo                                   Gly Gly

Mo Gly Gly Gly Trp Gly Gln Pro His Gly Gly Ser Trp Gly Gln Pro His 76
Bo                                                Gly

Mo Gly Gly Ser Trp Gly Gln Pro His Gly Gly Gly Trp Gly Gln --- --- 90
Bo              Gly                                         Pro His

Mo --- --- --- --- --- --- Gly Gly Gly Thr His Asn Gln Trp Asn Lys 100
Bo Gly Gly Gly Gly Trp Gly Gln                           Gly

Mo Pro Ser Lys Pro Lys Thr Asn Leu Lys His Val Ala Gly Ala Ala Ala 116
Bo                                      Met

Mo Ala Gly Ala Val Val Gly Gly Leu Gly Gly Tyr Met Leu Gly Ser Ala 132
Bo

Mo Met Ser Arg Pro Met Ile His Phe Gly Asn Asp Trp Glu Asp Arg Tyr 148
Bo              Leu              Ser      Tyr

Mo Tyr Arg Glu Asn Met Tyr Arg Tyr Pro Asn Gln Val Tyr Tyr Arg Pro 164
Bo              His

Mo Val Asp Gln Tyr Ser Asn Gln Asn Asn Phe Val His Asp Cys Val Asn 180
Bo

Mo Ile Thr Ile Lys Gln His Thr Val Thr Thr Thr Thr Lys Gly Glu Asn 200
Bo             Val      Glu

Mo Phe Thr Glu Thr Asp Val Lys Met Met Glu Arg Val Val Glu Gln Met 212
Bo                                Ile

Mo Cys Val Thr Gln Tyr Gln Lys Glu Ser Gln Ala Tyr Tyr Asp Gly Arg 228
Bo                                              Gln ---

Mo Arg Ser Ser Ser Thr Val Leu Phe Ser Ser Pro Pro Val Ile Leu Leu 244
Bo --- Gly Ala      Val Ile

Mo Ile Ser Phe Leu Ile Phe Leu Ile Val Gly 254
Bo
    (SEQ ID NO: 9)
    (SEQ ID NO: 7)

```

FIG. 3

Mo	Met	Ala	Asn	Leu	---	---	Gly	Tyr	Trp	Leu	Leu	Ala	Leu	Phe	Val	Thr	14
Sh		Val	Lys	Ser	His	Ile		Ser		Ile		Val				Ala	
Mo	Met	Trp	Thr	Asp	Val	Gly	Leu	Cys	Lys	Lys	Arg	Pro	Lys	Pro	Gly	Gly	30
Sh			Ser														
Mo	---	Trp	Asn	Thr	Gly	Gly	Ser	Arg	Tyr	Pro	Gly	Gln	Gly	Ser	Pro	Gly	45
Sh																	
Mo	Gly	Asn	Arg	Tyr	Pro	Pro	Gln	Gly	Gly	---	Thr	Trp	Gly	Gln	Pro	His	60
Sh										Gly	Gly						
Mo	Gly	Gly	Gly	Trp	Gly	Gln	Pro	His	Gly	Gly	Ser	Trp	Gly	Gln	Pro	His	76
Sh											Gly						
Mo	Gly	Gly	Ser	Trp	Gly	Gln	Pro	His	Gly	Gly	Gly	---	Trp	Gly	Gln	Gly	91
Sh												Gly					
Mo	Gly	Gly	Thr	His	Asn	Gln	Trp	Asn	Lys	Pro	Ser	Lys	Pro	Lys	Thr	Asn	107
Sh			Ser	---	His	Ser											
Mo	Leu	Lys	His	Val	Ala	Gly	Ala	Ala	Ala	Ala	Gly	Ala	Val	Val	Gly	Gly	123
Sh	Met																
Mo	Leu	Gly	Gly	Tyr	Met	Leu	Gly	Ser	Ala	Met	Ser	Arg	Pro	Met	Ile	His	139
Sh														Leu			
Mo	Phe	Gly	Asn	Asp	Trp	Glu	Asp	Arg	Tyr	Tyr	Arg	Glu	Asn	Met	Tyr	Arg	155
Sh					Tyr												
Mo	Tyr	Pro	Asn	Gln	Val	Tyr	Tyr	Arg	Pro	Val	Asp	Gln	Tyr	Ser	Asn	Gln	171
Sh																	
Mo	Asn	Asn	Phe	Val	His	Asp	Cys	Val	Asn	Ile	Thr	Ile	Lys	Gln	His	Thr	187
Sh												Val					
Mo	Val	Thr	Thr	Thr	Thr	Lys	Gly	Glu	Asn	Phe	Thr	Glu	Thr	Asp	Val	Lys	203
Sh															Ile		
Mo	Met	Met	Glu	Arg	Val	Val	Glu	Gln	Met	Cys	Val	Thr	Gln	Tyr	Gln	Lys	219
Sh	Ile										Ile				Arg		
Mo	Glu	Ser	Gln	Ala	Tyr	Tyr	Asp	Gly	Arg	Arg	Ser	Ser	Ser	Thr	Val	Leu	235
Sh							Gln	---	---		Gly	Ala		Val	Ile		
Mo	Phe	Ser	Ser	Pro	Pro	Val	Ile	Leu	Leu	Ile	Ser	Phe	Leu	Ile	Phe	Leu	251
Sh																	
Mo	Ile	Val	Gly														254
Sh																	

(SEQ ID NO: 10)
 (SEQ ID NO: 7)

FIG. 4

Generation of N-terminally located FLAG-tagged PrP constructs





FIG. 5



1. SPOX FLAG-MHM2 PrP
2. Tg(FLAG-MoPrP)FVB/Abi D7755

Survival of Tg(FLAG-MoPrP) after inoculation with Mouse RML prions




FIG. 6

RECOMBINANT CONSTRUCT ENCODING EPIOTOPE TAGGED PRP PROTEIN

CROSS REFERENCES

This application is a continuation application of Ser. No. 09/031,168, filed Feb. 26, 1998 now issued as U.S. Pat. No. 6,150,583, which is a divisional application of Ser. No. 08/660,626, filed Jun. 6, 1996, and now issued as U.S. Pat. No. 5,789,655, issued Aug. 4, 1998, which is a continuation-in-part application of Ser. No. 08/521,992, filed Aug. 31, 1995, now issued as U.S. Pat. No. 5,908,969, issued Jun. 1, 1999 which is a continuation-in-part of Ser. No. 08/509,261, filed Jul. 31, 1995, now issued as U.S. Pat. No. 5,763,740, issued Jun. 9, 1998, which is a continuation-in-part application of Ser. No. 08/242,188, filed May 13, 1994, and now issued as U.S. Pat. No. 5,565,186, issued Aug. 15, 1996, all of which are incorporated herein by reference in their entirety and to which applications we claim priority under 35 U.S.C. §120.

STATEMENT AS TO FEDERALLY SPONSORED RESEARCH

The United States Government may have certain rights in this application pursuant to Grant Nos. NS14069, NS22786, AG02132, AG08967, AG10770 and NS07219 awarded by the National Institutes of Health.

FIELD OF THE INVENTION

This invention relates generally to epitope-tagged proteins and to transgenic animals expressing such proteins. More specifically, this invention relates to epitope-tagged prion protein (PrP) genes, transgenic animals expressing epitope-tagged PrP genes, and assay methods for distinguishing between and isolating infectious and noninfectious prion proteins.

BACKGROUND OF THE INVENTION

Prions are infectious pathogens that cause central nervous system spongiform encephalopathies in humans and animals. Prions are distinct from bacteria, viruses and viroids. The predominant hypothesis at present is that no nucleic acid component is necessary for infectivity of prion protein. Further, a prion which infects one species of animal (e.g., a human) will not infect another (e.g., a mouse).

A major step in the study of prions and the diseases that they cause was the discovery and purification of a protein designated prion protein ("PrP") (Bolton et al. (1982) *Science* 218:1309-11; Prusiner et al. (1982) *Biochemistry* 21:6942-50; McKinley et al. (1983) *Cell* 35:57-62). Complete prion protein-encoding genes have since been cloned, sequenced and expressed in transgenic animals. PrP^C is encoded by a single-copy host gene (Basler et al. (1986) *Cell* 46:417-28) and is normally found at the outer surface of neurons. A leading hypothesis is that prion diseases result from conversion of PrP^C into a modified scrapie isoform called PrP^{Sc} during a post-translational process (Borchelt et al. (1990) *J. Cell Biol.* 110:743-752). It is likely that a fundamental event in the propagation of prions is the conformational transition of alpha-helices in PrP^C into beta-sheets in PrP^{Sc} (Pan et al. (1993) *Proc. Natl. Acad. Sci.* 90:10962-10966). Genetic evidence from transgenic mouse studies demonstrates the requirement for an additional component(s) referred to as protein X in this conversion (Telling et al. (1995) *Cell* 83:79-90).

It appears that PrP^{Sc} is necessary for both the transmission and pathogenesis of the transmissible neurodegenerative

diseases of animals and humans (see, Prusiner (1991) *Science* 252:1515-1522). The most common prion diseases of animals are scrapie of sheep and goats and bovine spongiform encephalopathy (BSE) of cattle (Wilesmith & Wells (1991) *Microbiol. Immunol.* 172:21-38). Four prion diseases of humans have been identified: (1) kuru, (2) Creutzfeldt-Jakob Disease (CJD), (3) Gerstmann-Strassler-Scheinker Disease (GSS), and (4) fatal familial insomnia (FFI) (Gajdusek (1977) *Science* 197:943-960; Medori et al. (1992) *N. Engl. J. Med.* 326:444-449). The presentation of human prion diseases as sporadic, genetic and infectious illnesses initially posed a conundrum which has been explained by the cellular genetic origin of PrP.

Most CJD cases are sporadic, but about 10-15% are inherited as autosomal dominant disorders that are caused by mutations in the human PrP gene (Hsiao et al. (1990) *Neurology* 40:1820-1827; Goldfarb et al. (1992) *Science* 258:806-808); Kitamoto et al. (1994) *Proc. R. Soc. Lond.* 343:391-398). Iatrogenic CJD has been caused by human growth hormone derived from cadaveric pituitaries as well as dura mater grafts (Brown et al. (1992) *Lancet* 340:24-27). attempts to link CJD to an infectious source such as the consumption of scrapie infected sheep meat, none has been identified to date (Harries-Jones et al. (1988) *J. Neurol. Neurosurg. Psychiatry* 51:1113-1119) except in cases of iatrogenically induced disease. On the other hand, kuru, which for many decades devastated the Fore and neighboring tribes of the New Guinea highlands, is believed to have been spread by infection during ritualistic cannibalism (Alpers (1979) *Slow Transmissible Diseases of the Nervous System*, Vol. 1, S. B. Prusiner and W. J. Hadlow, eds. (New York: Academic Press), pp. 66-90).

The initial transmission of CJD to experimental primates has a rich history beginning with William Hadlow's recognition of the similarity between kuru and scrapie. In 1959, Hadlow suggested that extracts prepared from patients dying of kuru be inoculated into non-human primates and that the animals be observed for disease that was predicted to occur after a prolonged incubation period (Hadlow (1959) *Lancet* 2:289-290). Seven years later, Gajdusek, Gibbs and Alpers demonstrated the transmissibility of kuru to chimpanzees after incubation periods ranging from 18 to 21 months (Gajdusek et al. (1966) *Nature* 209:794-796). The similarity of the neuropathology of kuru with that of CJD (Klatzo et al. (1959) *Lab Invest.* 8:799-847) prompted similar experiments with chimpanzees and transmissions of disease were reported in 1968 (Gibbs, Jr. et al. (1968) *Science* 161:388-389). Over the last 25 years, about 300 cases of CJD, kuru and GSS have been transmitted to a variety of apes and monkeys.

The expense, scarcity and often perceived inhumanity of such experiments have restricted this work and thus limited the accumulation of knowledge. While the most reliable transmission data has been said to emanate from studies using non-human primates, some cases of human prion disease have been transmitted to rodents but apparently with less regularity (Gibbs, Jr. et al. (1979) *Slow Transmissible Diseases of the Nervous System*, Vol. 2, S. B. Prusiner and W. J. Hadlow, eds. (New York: Academic Press), pp. 87-110; Tateishi et al. (1992) *Prion Diseases of Humans and Animals*, Prusiner et al., eds. (London: Ellis Horwood), pp. 129-134).

The infrequent transmission of human prion disease to rodents has been cited as an example of the "species barrier" first described by Pattison in his studies of passaging the scrapie agent between sheep and rodents (Pattison (1965) *NINDB Monograph* 2, D. C. Gajdusek, C. J. Gibbs Jr. and

M. P. Alpers, eds. (Washington, D.C.: U.S. Government Printing), pp. 249–257). In those investigations, the initial passage of prions from one species to another was associated with a prolonged incubation time with only a few animals developing illness. Subsequent passage in the same species was characterized by all the animals becoming ill after greatly shortened incubation times.

The molecular basis for the species barrier between Syrian hamster (SHa) and mouse (Mo) was shown to reside in the species-specific differences in the sequence of the PrP (Scott et al. (1989) *Cell* 59:847–857). Mouse PrP (MoPrP) differs from Syrian hamster PrP (SHaPrP) at 16 positions out of 254 amino acid residues (Basler et al. (1986), *Cell* supra; Lochter et al. (1986) *Proc. Natl. Acad. Sci. USA* 83:6372–6376). Transgenic mice expressing SHaPrP [Tg (SHaPrP)] had abbreviated incubation times when inoculated with SHa prions. When similar studies were performed with mice expressing the human, or ovine PrP transgenes, the species barrier was not abrogated, i.e., the percentage of animals which became infected were unacceptably low and the incubation times were unacceptably long (Telling et al. (1994) *Proc. Natl. Acad. Sci.* 91:9936–9940; Telling et al. (1995) *Cell* 83:79–90). Thus, it was not possible to render non-human animals such as mice, susceptible to infection by human prions.

Purification of PrP^{Sc} has been facilitated by its relative resistance to proteolytic degradation and insolubility in non-denaturing detergents (Bolton et al. (1982) supra; Prusiner et al. (1982) supra). Purification of PrP^C has been more problematic. Immunoaffinity chromatography purification of PrP^C yielded only small amounts of protein. Improved purification of PrP^C has been accomplished by a multi-step purification procedure involving detergent extraction and separation by immobilized Cu²⁺ ion affinity chromatography followed by cation-exchange chromatography and sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) (Pan et al. (1992) *Protein Sci.* 1:136–144).

The production of monoclonal antibodies against PrP^C and PrP^{Sc} has been particularly difficult. In the case of mouse PrP, MoPrP is recognized as self, precluding the production of anti-MoPrP antibodies in animals immunized with MoPrP.

There is an urgent need to develop diagnostics and therapeutics for PrP^{Sc}-mediated diseases such as CJD. Although many lines of evidence support the idea that PrP^C is converted to the infectious PrP^{Sc} isoform, greater understanding of the conditions under which scrapie infectivity is generated de novo is needed to develop compounds able to inhibit the generation of PrP^{Sc}. Compounds able to inhibit the in vitro conversion of PrP^C to PrP^{Sc} could be useful for the treatment and prevention of prion-mediated diseases in animal and human subjects at risk. Improved methods for monitoring the conversion of PrP from the alpha-helical conformation of PrP^C to the beta-sheet conformation of the infectious PrP^{Sc} isoform would be useful in developing assays for such compounds.

SUMMARY OF THE INVENTION

Nucleotides encoding a strong epitope tag are operatively placed in a nucleotide sequence encoding a protein which normally has two or more conformational shapes. Depending on the conformational shape assumed by the expressed protein, the tag will or will not be exposed thereby making it possible to differentiate between conformational shapes via an antibody which binds to the epitope. An aspect of the

invention features a recombinant nucleic acid construct comprising a nucleic acid sequence encoding an amino acid sequence comprising a biologically active protein or protein fragment connected, preferably directly, to a heterologous epitope domain. The expressed amino acid sequence (i.e., the epitope-tagged protein) preferably retains the biological activity of the corresponding natural (e.g., untagged) protein or protein fragment. The tag may be used in connection with a protein which has two or more different conformational shapes, such that the epitope tag is relatively more exposed in one conformational shape relative to another conformational shape.

One aspect of the invention is a transgenic animal such as a mouse which has incorporated into its genome a first DNA sequence encoding a protein which when expressed assumes two or more different conformational shapes. The first DNA sequence has a second DNA sequence encoding an epitope tag connected to it. The second sequence is preferably positioned relative to the first sequence such that the exposure of the tag after expression changes with the different conformational shapes assumed by the protein expressed by the first sequence. The first DNA sequence is preferably an exogenous sequence which encodes a protein such as PrP which protein causes a disease in one conformational shape but not another. Thus by correctly positioning the second sequence encoding the tag relative to the first sequence, it is possible to quickly and easily assay a sample from the animal and determine which conformation the protein has assumed.

Transgenic mammals comprising a tagged transgene are preferably selected from the group consisting of *Mus*, *Rattus*, *Oryctolagus* and *Mesocricetus*. Transgenic animals expressing high levels of the tagged transgene may be obtained, for example, by over-expression of the transgene with an enhanced promoter and/or with high copy numbers of the transgene.

In a specific embodiment, the invention features a transgenic mammal having an epitope-tagged PrP gene. The PrP gene may be a natural, synthetic, or chimeric PrP gene. In specific embodiments, the transgenic animals have an epitope-tagged chimeric PrP gene which renders the transgenic animals susceptible to infection with a prion which generally only infects a genetically diverse or distinct animal. A chimeric PrP gene is a gene which includes a portion of a gene of a genetically diverse animal. When the transgenic animal is a one of *Mus*, *Rattus*, *Oryctolagus*, or *Mesocricetus*, the genetically diverse or distinct animal is selected from the group consisting of *Bos*, *Ovis*, *Sus*, and *Homo*. A preferred transgenic animal is a mouse expressing an epitope-tagged chimeric PrP in which a segment of mouse (Mo) PrP is replaced with the corresponding human (Hu) PrP sequence.

The transgenic animal may be heterozygous or homozygous for an ablated or disrupted endogenous PrP gene; in a preferred embodiment, the transgenic animal is homozygous for an ablated endogenous PrP gene.

In a preferred embodiment of the invention, the epitope-tagged protein is a natural, synthetic, or chimeric prion protein (PrP). PrP may be tagged with a variety of natural or artificial heterologous epitopes known in the art, including artificial epitopes such as FLAG, Strep, or poly-histidine peptides. FLAG peptides include the sequence Asp-Tyr-Lys-Asp-Asp-Asp-Asp-Lys (SEQ ID NO:1) or Asp-Tyr-Lys-Asp-Glu-Asp-Asp-Lys (SEQ ID NO:2). The Strep epitope has the sequence Ala-Trp-Arg-His-Pro-Gln-Phe-Gly-Gly (SEQ ID NO:3). Another commonly used artificial epitope

is a poly-His sequence having six histidine residues (His-His-His-His-His-His) (SEQ ID NO:4). Naturally-occurring epitopes include the influenza virus hemagglutinin sequence Tyr-Pro-Tyr-Asp-Val-Pro-Asp-Tyr-Ala-Ile-Glu-Gly-Arg (SEQ ID NO:11) recognized by the monoclonal antibody 12CA5 (Murray et al. (1995) *Anal. Biochem.* 229:170–179) and the eleven amino acid sequence from human c-myc recognized by the monoclonal antibody 9E10 (Glu-Gln-Lys-Leu-Leu-Ser-Glu-Glu-Asp-Leu-Asn) (SEQ ID NO:12) (Manstein et al. (1995) *Gene* 162:129–134). Another useful epitope is the tripeptide Glu-Glu-Phe which is recognized by the monoclonal antibody YL 1/2 against α -tubulin. This tripeptide has been used as an affinity tag for the purification of recombinant proteins (Stammers et al. (1991) *FEBS Lett.* 283:298–302).

In a particularly preferred embodiment, the epitope tagged PrP molecule has an artificial FLAG epitope inserted after codon 22, e.g., the first codon of the FLAG epitope begins at codon 23 of a nucleotide sequence encoding a FLAG-tagged PrP protein. The FLAG-tagged PrP molecule retains all of the biological activity of the natural PrP molecule. Specifically, the FLAG-tagged PrP protein retains the ability to support prion propagation.

The invention further includes cells, e.g., omnipotent and pluripotent cells, and immortalized cell lines expressing the epitope-tagged protein construct, as well as transgenic animals having a gene encoding an epitope-tagged protein integrated into their genome.

In another aspect, the invention features a method for distinguishing between the conformational shapes of a protein having a first and second conformation shape, comprising the steps of: (a) generating a recombinant nucleic acid construct comprising a nucleic acid sequence encoding an amino acid sequence comprising a protein fragment tagged with a heterologous epitope; b) transfecting a cell or organism with the tagged protein construct; c) expressing the tagged protein. The epitope tag is positioned relative to the protein sequence such that the epitope is exposed on the surface of the tagged protein to a greater degree when the protein is in a first conformational shape relative to the degree of exposure of the epitope when the protein is in a second conformational shape. In one embodiment, the conformational shapes of the protein can be distinguished by detecting the presence or absence of the epitope. Multiple different tags can be used if the protein assumes multiple conformations, making it possible to distinguish the conformations via detection of the presence or absence of a series of tags. In another embodiment, the conformational shapes of a protein are distinguished by relatively greater exposure of the epitope tag in one conformational shape than in other conformational shapes. Preferably, the exposure of an epitope tag is 20–100% greater in one conformational shape relative to the second conformational shape; more preferably, the relative exposure is 50–100% greater; most preferably, the relative exposure is 75–100% greater.

In one embodiment, the epitope-tagged protein is PrP, and the epitope tag is placed such that it is exposed on the surface of the expressed prion protein when it is in the noninfectious alpha-helical PrP^C isoform, but the epitope tag is exposed on the surface of the infectious beta-sheet PrP^{Sc} isoform.

In another aspect, the invention features a method of isolating PrP by a) generating a recombinant nucleic acid construct comprising a nucleic acid sequence encoding a prion protein having a heterologous epitope domain; b) transfecting a cell or organism with the tagged PrP construct;

c) expressing the construct to produce epitope-tagged PrP, where the epitope tag is placed such that it is exposed on the surface of the desired PrP isoform; and d) purifying PrP by immunoaffinity chromatography using an anti-epitope tag antibody. In a specific embodiment, the method of isolating PrP includes an additional step of enriching for PrP prior to purification. This method can be used to isolate, separate and identify either PrP^C and/or PrP^{Sc}.

In another aspect, the invention features an assay method for detecting infectious prions by a) generating a transgenic animal comprised of an epitope-tagged PrP gene where the epitope tag is relatively more exposed on the surface of the expressed PrP molecule when the molecule has the PrP^{Sc} conformation than when the molecule is in the PrP^C conformation; b) inoculating the transgenic animal with material suspected of containing infectious prions; and c) detecting the increased presence of epitope-tagged PrP. Detection of increased levels of epitope-tagged PrP results from increased levels of PrP in the infectious PrP^{Sc} conformation, thus indicating the presence of infectious PrP particles in the inoculating material. In one preferred embodiment, the transgenic animal expresses a bovine-mouse MBov2M chimeric PrP gene and is inoculated with material from infected cattle. In another preferred embodiment, the transgenic animal expresses a chimeric human-mouse MHu2M PrP molecule and is inoculated from material from an infected human.

One object of the invention is to provide a transgenic animal producing large quantities of an epitope-tagged protein or protein fragment which is easily purified via immunoaffinity chromatography using an epitope-specific antibody. This is particularly useful where the protein is difficult to purify in sufficient quantities and/or attempts to produce antibodies specific to the protein and its conformation isoforms have been unsuccessful, e.g., PrP^C and PrP^{Sc}. Additionally, the invention allows a simplified one-step enrichment of PrP^C and/or PrP^{Sc}, which can be followed by a variety of procedures including immunodetection.

Another object is to provide a transgenic animal expressing elevated levels of a tagged protein or protein fragment obtained with an enhanced promoter or a high copy number of a tagged transgene.

Another object is to provide a method for distinguishing conformational changes in a protein, e.g., distinguishing between the isoforms of PrP^C and PrP^{Sc}.

Another object is to provide a gene tagged with a heterologous epitope.

Another object of the invention is to provide a transgenic host mammal (which is small, e.g., less than 1 kg when full grown, and inexpensive to maintain) such as a mouse, rat or hamster which includes an exogenous or chimeric PrP gene, including all or a portion of a PrP gene from another animal, (which is large, greater than 2 kg when full grown, and expensive to maintain) such as a human, cow, pig, sheep, cat or dog, and having a artificial epitope tag domain.

Another object of the invention is to provide a transgenic host animal which includes elevated levels of expression of a tagged PrP gene of a genetically diverse animal wherein the elevated levels of expression are obtained by the inclusion of a high copy number of the tagged PrP gene of the genetically diverse mammal and/or fusing an enhanced promoter to the PrP gene of the genetically diverse animal.

One advantage of the method of the invention is the production of elevated levels of readily isolatable PrP^C and PrP^{Sc}.

Another object is to provide a transgenic animal assay which animal, on inoculation, develops PrP^{Sc} which is detectable via an epitope tag as distinguished from PrP^C.

These and other objects, advantages and features of the present invention will become apparent to those persons skilled in the art upon reading the details of the compositions, composition components, methods and method steps of the invention as set forth below.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 shows the amino acid sequence of mouse PrP along with specific differences between mouse PrP and human PrP (SEQ ID NO:8).

FIG. 2 shows the amino acid sequence of mouse PrP (SEQ ID NO:7) along with specific differences between mouse PrP and bovine PrP (SEQ ID NO:9).

FIG. 3 shows the amino acid sequence of mouse PrP (SEQ ID NO:7) along with specific differences between mouse PrP and sheep PrP (SEQ ID NO:10).

FIG. 4 is the PrP amino acid sequence showing insertion of a StuI recognition site upstream of the signal peptidase cleavage site at amino acid 2 (SEQ ID NO:14).

FIG. 5 is the amino acid sequence of tagged PrP having the FLAG epitope inserted at amino acid 2 (SEQ ID NO:15).

FIG. 6 is a chart showing the survival of Tg(FLAG-MoPrP) relative to non-transgenic mice ("Non-Tg mice") and a transgenic mouse line which overexpresses normal mouse PrP ("Tg(MoPrP-A)") after inoculation with mouse RML prions.

DETAILED DESCRIPTION

Before the present artificial epitope-tagged gene, assay methodology, and transgenic animals used in the assay are described, it is to be understood that this invention is not limited to particular assay methods, epitope-tagged and artificial genes, or transgenic animals described, as such methods, genes, and animals may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference to disclose and describe the methods and/or materials in connection with which the publications are cited. The publications discussed above are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

Definitions

The term "artificial" as used with "artificial gene" or "artificial epitope" and the like, refers to a non-naturally occurring material, e.g., a nucleotide sequence manufactured by human intervention, e.g., by fusing natural sequences together or chemically synthesizing sequences in isolation. Further, the term is intended to encompass a natural sequence which may be isolated from a naturally occurring genome and then connected, artificially, with an sequence (either a natural or artificial sequence) with which it is not naturally connected, e.g., a natural HIV virus epitope

connected directly to a natural PrP sequence is expressed by an "artificial gene."

The term "transgene" or "transgenic element" refers to an artificially introduced, chromosomally integrated nucleic acid sequence heterologous to the genome of the host animal in which the nucleic acid sequence is present.

The term "transgenic animal" means a non-human mammalian animal having a transgenic element integrated in its genome.

The term "prion" means an infectious particle known to cause diseases (spongiform encephalopathies) in humans and animals. The term "prion" is a contraction of the words "protein" and "infection" and the particles are comprised largely if not exclusively of PrP^{Sc} molecules encoded by a PrP gene. Prions are distinct from bacteria, viruses and viroids. Known prions include those which infect animals to cause scrapie, a transmissible, degenerative disease of the nervous system of sheep and goats as well as bovine spongiform encephalopathies (BSE) or mad cow disease and feline spongiform encephalopathies of cats. Four prion diseases known to affect humans are (1) kuru, (2) Creutzfeldt-Jakob Disease (CJD), (3) Gerstmann-Strassler-Scheinker Disease (GSS), and (4) fatal familial insomnia (FFI). As used herein prion includes all forms of prions causing all or any of these diseases or others in any animals used—and in particular in humans and in domesticated farm animals.

The term "PrP gene", "prion protein gene" or "PrP sequence" are used interchangeably herein to describe genetic material which expresses any PrP proteins, for example those shown in FIGS. 1-3. There are a number of known variants to the human PrP gene. Further, there are known polymorphisms in the human, sheep, and bovine PrP gene. The following is a list of such variants:

Pathogenic human mutations	Human Polymorphisms	Sheep Polymorphisms	Bovine Polymorphisms
2 octarepeat insert	Codon 129 Met/Val	Codon 171 Arg/Glu	5 or 6 octarepeat
4 octarepeat insert	Codon 219 Glu/Lys	Codon 136 Ala/Val	
5 octarepeat insert			
6 octarepeat insert			
7 octarepeat insert			
8 octarepeat insert			
9 octarepeat insert			
Codon 102 Pro-Leu			
Codon 105 Pro-Leu			
Codon 117 Ala-Val			
Codon 145 Stop			
Codon 178 Asp-Asn			
Codon 180 Val-Ile			
Codon 198 Phe-Ser			
Codon 200 Glu-Lys			
Codon 210 Val-Ile			
Codon 217 Asn-Arg			

-continued

Pathogenic human mutations	Human Polymorphisms	Sheep Polymorphisms	Bovine Polymorphisms
Codon 232			
Met-Ala			

The PrP gene can be a naturally-occurring PrP gene from any animal described herein and any and all polymorphisms and mutations thereof, it being recognized that the terms include other such PrP genes that are yet to be discovered. The term "PrP gene" generally includes any gene of any species which encodes any form of a prion protein. Some commonly known PrP sequences are described in Gabriel et al., *Proc. Natl. Acad. Sci. USA* 89:9097-9101 (1992) which is incorporated herein by reference to disclose and describe such sequences. Besides naturally-occurring PrP genes, the term "PrP gene" further encompasses artificial, synthetic, and chimeric Prp genes.

The term "chimeric PrP gene" is used herein to encompass recombinantly constructed genes which when included in the genome of a host animal (e.g., a mouse) will render the mammal susceptible to infection from prions which naturally only infect a genetically diverse test mammal, e.g., human, bovine or ovine. A chimeric PrP will accomplish this effect in an animal which includes an operative endogenous PrP gene and allow the animal to show symptoms within 200±50 days after inoculation. In general, a chimeric gene will include the codon sequence of the PrP gene of the mammal being genetically altered with one or more (but not all, and generally less than 40) codons of the natural sequence being replaced with a different codon—preferably a corresponding codon of a genetically diverse mammal (such as a human). The genetically altered mammal being used to assay samples for prions which only infect the genetically diverse mammal. Examples of artificial genes are mouse PrP genes encoding the sequence as shown in FIGS. 1, 2 and 3 with one or more different replacement codons selected from the codons shown in these Figures for humans, cows and sheep replacing mouse codons at the same position, with the proviso that not all the mouse codons are replaced with differing human, cow or sheep codons. The chimeric PrP genes of the invention can include not only codons of genetically diverse animals but may include codons and codon sequences not associated with any native PrP gene but which, when inserted into an animal render the animal susceptible to infection with prions which would normally only infect a genetically diverse animal. The chimeric PrP genes can also include deletions of codons in the naturally-occurring gene of the host animal genome.

The terms "heterologous epitope domain" or "heterologous epitope tag" are used interchangeably to mean a peptide which does not naturally occur as part of the protein or protein fragment. The heterologous epitope may be a naturally-occurring or artificial peptide. Heterologous epitope tags are distinguished from naturally occurring antigenic PrP sequences to which monoclonal antibodies such as 3F4 and 13AS have been raised. The heterologous epitope tag does not necessarily have to be antigenic, e.g., generate a specific antibody. The epitope tag includes peptides bound by specific ligands. For example, an Arg-Gly-Asp sequence may be inserted into the PrP protein which binds integrins (Ruoslahti & Eierschbacher (1987) *Science* 238:491-497).

The term "heterologous epitope-tagged gene" or "tagged gene" are used to mean an artificially constructed nucleotide

sequence comprising a gene (i.e., a first nucleotide sequence) encoding an amino acid sequence comprising a biologically active protein or protein fragment of interest and a second nucleotide sequences encoding an epitope tag which does not naturally occur as part of the protein or protein fragment, e.g., heterologous. The sequence encoding the tag may be placed 5', 3', and/or within the sequence encoding the protein of interest. Expression of the tagged gene results in production of a protein or protein fragment having one or more heterologous epitope domain(s).

The terms "epitope-tagged PrP gene" or "tagged PrP gene", and the like are used interchangeably herein to mean an artificially constructed gene containing the nucleotide sequence encoding the PrP protein of an animal such as a mouse, human, cow, or sheep and additionally containing nucleotide sequences encoding one or more epitope tags inserted into the PrP gene. Expression of the tagged PrP gene results in production of PrP molecule having one or more artificial epitope domain(s). In one specific example the tagged PrP gene is comprised of the nucleotide sequence encoding mouse PrP with a nucleotide sequence encoding an artificial FLAG epitope inserted at codon 22. In another specific embodiment, the tagged-PrP gene comprises the nucleotide sequence encoding chimeric mouse-hamster PrP (MHM2) with the nucleotide sequence encoding a FLAG epitope inserted at codon 22 of the MHM2 PrP gene. The nucleotide sequence also contains the epitope derived from hamster PrP recognized by the monoclonal antibody 3F4 (Scott et al. (1992) *Protein Sci.* 1:986-997). The resulting tagged PrP molecule can be readily purified through immunoaffinity methodologies with the use of antibodies to the epitope tag(s).

The terms "epitope-tagged chimeric PrP gene" or "tagged chimeric PrP gene", and the like are used interchangeably herein to mean an artificially constructed PrP gene containing the nucleotide sequence encoding the PrP protein of a host animal such as a mouse with one or more of the codons being replaced with corresponding codons from a genetically diverse test animal such as a human, cow or sheep, and having a nucleotide sequence encoding a heterologous epitope tag inserted into the PrP gene, such that expression of the tagged PrP gene results in production of a PrP protein having one or more heterologous epitope domain(s). In one specific example the tagged PrP gene is comprised of the starting and terminating sequence (i.e., N- and C-terminal codons) of a PrP gene of a mammal of a host species (e.g. a mouse), a nucleotide sequence of a corresponding portion of a PrP gene of a test mammal of a second species (e.g. a human), and a nucleotide sequence encoding a FLAG epitope at codon 22 (FIG. 5). When inserted into the genome of a mammal of the host species, the tagged PrP gene will render the mammal susceptible to infection with prions which normally infect only mammals of the second, genetically diverse, species, i.e. on inoculation the host animal will show symptoms of prion disease within 200±50 days. A preferred tagged PrP gene is MHu2M which contains the starting and terminating sequence of a mouse PrP gene and a non-terminal sequence region which is replaced with a corresponding human sequence which differs from a mouse PrP gene in a manner such that the protein expressed thereby differs at nine residues (MHu2M).

The term "genetic material related to prions" is intended to cover any genetic material which effects the ability of an animal to become infected with prions. Thus, the term encompasses any "PrP gene", "chimeric PrP gene", or "ablated PrP gene" which terms are defined herein as well as modification of such which effect the ability of an animal to become infected with prions.

11

The terms “host animal,” “host non-human mammal,” and the like are used to describe animals which will have their genome genetically and artificially manipulated so as to include genetic material which is not naturally present within the animal. For example, host animals include mice, hamsters and rats which have their PrP gene disrupted or altered by the insertion of a tagged artificial PrP gene of another animal or by the insertion of a tagged native PrP gene of a genetically diverse test animal.

The terms “ablated prion protein gene”, “disrupted PrP gene”, and the like are used interchangeably herein to mean an endogenous prion protein gene which has been altered (e.g., add and/or remove nucleotides) in a manner so as to render the gene inoperative. Examples of non-functional prion protein genes and methods of making such are disclosed in Büeler et al. (1992) *Nature* 356:577–582, which is incorporated herein by reference. One (heterozygous) or preferably both (homozygous) alleles of the genes are disrupted.

The terms “retaining the biological activity,” “the biological activity of the naturally occurring protein or protein fragment”, and the like, mean that the tagged protein or tagged protein fragment retains at least part of and preferably all of the characteristic biological activities and specificities of the unmodified, e.g., untagged, protein or protein fragment. For example, the FLAG-tagged chimeric Syrian hamster-mouse PrP (FLAG-MHM2PrP) retains the ability to support prion propagation *in vivo* when expressed in transgenic mice infected with Syrian hamster prions.

The terms “genetically diverse animal” and “genetically diverse mammal” are used to describe an animal which includes a native PrP codon sequence of the host animal which differs from the genetically diverse test animal by 17 or more codons, preferably 20 or more codons, and most preferably 28–40 codons. Thus, a mouse PrP gene is genetically diverse with respect to the PrP gene of a human cow or sheep, but is not genetically diverse with respect to the PrP gene of a hamster.

The terms “susceptible to infection” and “susceptible to infection by prions” and the like are used interchangeably herein to describe a transgenic animal of the invention which has an 80% or greater, preferably 98% or greater, and most preferably a 100% chance of developing a disease if inoculated with prions which would not normally infect a genetically diverse animal. Further, an animal “susceptible to infection” will develop symptoms of prion disease with 200±50 days or less after inoculation with prions which they are susceptible to being infected with.

The terms are used to describe a transgenic animal of the invention such as a transgenic mouse Tg(MHu2M) which, without the chimeric PrP gene, would rarely be susceptible to infection with a human or bovine prion (less than 20% chance of infection), but with the chimeric gene is susceptible to infection with human or bovine prions (80% to 100% chance of infection) and will show symptoms within 250 days or less after inoculation.

The term “incubation time” means the time from inoculation of an animal with a prion until the time when the animal first develops detectable symptoms of disease resulting from the infection. A reduced incubation time is one year or less, preferable about 200 days±50 days or less, more preferably about 50 days±20 days or less. Generally, in connection with the present invention, “incubation time” means the time from inoculation of an animal with any substance which causes a conformational change in a natural protein until that conformational change takes place in a detectable amount, e.g., detecting the exposed tag.

12

Epitope Tags

The invention includes epitope-tagged transgenes which are recombinant nucleic acid constructs encoding an amino acid sequence. The construct comprises a first sequence encoding biologically active protein or biologically active fragment and a second sequence coding for a tag with a heterologous epitope domain. The invention also includes transgenic animals expressing an epitope-tagged transgene. A variety of epitopes may be used to tag a protein, so long as the epitope (1) is heterologous to the naturally-occurring protein, and (2) the epitope-tagged protein retains at least part and preferably all of the biological activity of the unmodified protein. Such epitopes may be naturally-occurring amino acid sequences found in nature, artificially constructed sequences, or modified natural sequences. Recently, a variety of artificial epitope sequences have been described that have been shown to be useful for tagging and detecting recombinant proteins. One such tag, the eight amino acid FLAG marker peptide (Asp-Tyr-Lys-Asp-Asp-Asp-Lys) (SEQ ID NO:1), has a number of features which make it particularly useful for not only detection but also affinity purification of recombinant proteins (Brewer (1991) *Bioprocess Technol.* 2:239–266; Kunz (1992) *J. Biol. Chem.* 267:9101–9106). Inclusion of the FLAG epitope in recombinant proteins avoids the necessity for the development of a specialized scheme or functional assay for protein purification and circumvents need to raise antibodies against the tagged protein, the first four amino acids of this sequence comprising the antigenic site for α -FLAG M1 and M2 monoclonal antibodies. The small octapeptide has a high degree of hydrophilicity, thus maximizing accessibility to α -FLAG M1 and M2 monoclonal antibodies. A particularly useful feature is the calcium-dependent binding of the α -FLAG M1 monoclonal antibody to recombinant proteins containing the FLAG peptide. Removal of the Ca^{2+} by chelation with EDTA allows for efficient immunoaffinity purification without denaturation. A further advantage of the FLAG system is that it allows cleavage of the FLAG peptide from purified protein since the tag contains the rare five amino acid recognition sequence for enterokinase. The anti-FLAG M1 antibody requires an N-terminal FLAG sequence. A second anti-FLAG monoclonal antibody (anti-FLAG M2) has been employed in immunoaffinity purification of N-terminal Met-FLAG and C-terminal FLAG fusion proteins (Brizzard et al. (1994) *Biotechniques* 16:730–735). This antibody has, however, been found to cross-react with a splicing isoform of Mg^{2+} dependent protein phosphatase beta (MPP beta) which contains a sequence motif with five out of eight amino acid residues identical to the FLAG peptide (Schafer (1995) *Biochem. Biophys. Res. Commun.* 207:708–714). Binding of an anti-FLAG M2 monoclonal antibody to the FLAG epitope is not calcium-dependent, but bound fusion proteins can be eluted by competition with FLAG peptide.

Additional artificial epitope tags include an improved FLAG tag having the sequence Asp-Tyr-Lys-Asp-Glu-Asp-Asp-Lys (SEQ ID NO:2), a nine amino acid peptide sequence Ala-Trp-Arg-His-Pro-Gln-Phe-Gly-Gly (SEQ ID NO:3) referred to as the “Strep tag” (Schmidt (1994) *J. Chromatography* 676:337–345), poly-histidine sequences, e.g., a poly-His of six residues which is sufficient for binding to IMAC beads, an eleven amino acid sequence from human c-myc recognized by monoclonal antibody 9E10, or an epitope represented by the sequence Tyr-Pro-Tyr-Asp-Val-Pro-Asp-Tyr-Ala-Ile-Glu-Gly-Arg (SEQ ID NO:11) derived from an influenza virus hemagglutinin (HA) subtype, recognized by the monoclonal antibody 12CA5. Also, the

Glu-Glu-Phe sequence recognized by the anti- α -tubulin monoclonal antibody YL1/2 has been used as an affinity tag for purification of recombinant proteins (Stammers et al. (1991) FEBS Lett. 283:298–302).

The present invention features transgenic animals having a transgene encoding a tagged protein or protein fragment. A variety of natural, modified, or artificial epitope tags may be used so long as insertion into the gene construct does not completely interfere with the biological activity of the encoded protein.

Natural and Chimeric PrP Gene

The present invention may be used to generate transgenic animals carrying a DNA construct encoding an epitope-tagged naturally occurring or chimeric PrP gene. Examples of naturally occurring PrP proteins are shown in FIGS. 1–3. PrP genes from any animal of interest may be epitope-tagged so long as the resulting tagged PrP molecule retains the biological activity of the natural protein, e.g., the ability of tagged PrP^C to be converted into PrP^{Sc} and to propagate prions.

Chimeric PrP transgenes are described in U.S. Pat. No. 5,565,186, issued Oct. 15, 1996, U.S. Pat. No. 5,763,740 issued Jun. 9, 1998, and U.S. Pat. No. 5,908,969 issued Jun. 1, 1999, all of which are herein specifically incorporated by reference, including the chimeric Syrian hamster/mouse (SHa/Mo) transgene MH2M which carries 5 amino acid substitutions found in SHaPrP lying between codons 94 and 188 (Scott et al. (1993) Cell 73:979–988), and the chimeric human/mouse PrP gene, MHu2M, in which the same region of the mouse gene is replaced by the corresponding human sequence which differs from the mouse PrP at 9 codons.

Mice expressing the MHu2M chimeric PrP transgene have been shown to be susceptible to human prions after abbreviated incubation times. That is, transgenic mice carrying the MHu2M gene [Tg(MHu2M)] will, after inoculation with human prions, develop disease symptoms attributed to the prions within about 180 days. Further, 80% or more of the transgenic mice inoculated with human prions will develop symptoms of the disease. Thus, PrP transgenic animals provide an excellent system for assessing prion infections.

Transgenic Animals

The invention features transgenic animals having a transgene encoding an amino acid sequence comprising a biologically active protein fragment and a heterologous epitope domain. Transgenic animals having a tagged protein transgene are generated by introducing the DNA constructs encoding the desired tagged protein into the germline DNA of a host animal. Several methods for generating transgenic animals are known in the art, see for example, Gordon et al. (1980) Proc. Natl. Acad. Sci. 77:7380–7384, herein specifically incorporated by reference for methods of genetically transforming embryos by microinjection of DNA). Introduction of the desired tagged protein sequences can also be accomplished by microinjection into a fertilized egg of the host animal; transformation of embryonic stem (ES) cells with the desired DNA and introduction of the transformed ES cells into host animal blastocysts; or embryonic transduction with a retroviral vector containing the desired transgene. Another method of generating transgenic animals is described in U.S. Pat. No. 5,487,992, which uses a positive-negative selector (PNS) vector for inserting a DNA sequence by homologous recombination into a target site in the host animal genome.

Generation of transgenic animals by microinjection techniques is well known in the art. To generate tagged protein transgenic mice, for example, a DNA fragment encoding the protein of interest into which the epitope tag has been inserted is prepared and microinjected into fertilized eggs of mice, followed by transfer of viable eggs into the oviducts of pseudopregnant mice (Hogan et al. (1986) *Manipulation of Mouse Embryos: A Laboratory Manual*, Cold Spring Harbor Laboratory, N. Y., herein specifically incorporated by reference for methods of generating transgenic animals).

In a preferred embodiment, transgenic animals are generated having a tagged PrP transgene. In a specific embodiment, the tagged PrP transgene is a PrP sequence tagged with the FLAG epitope at codon 22 (FIG. 5). Preferably, the transgenic animal is an animal which does not express the endogenous PrP gene. The transgenic mouse may be heterozygous or homozygous for modifications in one or both alleles resulting in effective deletion of the endogenous PrP gene (Δ PrP). Preferably, both alleles of the genes are disrupted. The endogenous prion protein gene may be altered in any manner (e.g., add and/or remove nucleotides) so as to render the gene inoperative. Examples of non-functional prion protein genes and methods of making such are disclosed in Büeler et al. (1992) Nature 356:577–582, which is incorporated herein by reference.

Expression of a Biologically Active Tagged PrP Protein

Previous efforts to generate a MoPrP^C specific antibody have been unsuccessful, likely resulting from recognition of MoPrP as self in immunized animals. This limitation has now been overcome by tagging PrP with the FLAG epitope tag. The FLAG system has the advantage of efficient, one-step immunoaffinity purification without denaturation using the anti-FLAG M1 monoclonal antibody which binds to proteins containing the FLAG peptide in a calcium-dependent manner. Recognition of FLAG fusion proteins by the anti-FLAG M1 monoclonal antibody relies on the location of the FLAG sequence at the N-terminus of the protein. Since PrP is processed in cells by the removal of a N-terminal signal peptide, the FLAG sequence was inserted distal to the signal peptidase cleavage site at amino acid residue 22 of PrP.

To construct FLAG-tagged PrP, a new recognition sequence for the restriction enzyme StuI was created upstream of the signal peptidase cleavage site which cuts PrP after amino acid 22. The unique StuI site was created by changing a T for an A at nucleotide position 57 of the MoPrP gene by the PCR-mediated mutagenesis described in Example 1 (FIG. 4). Variation at this nucleotide position do not change the predicted primary structure of PrP. Complementary oligonucleotide sequences [TTGGCCGCTTCT TGTCATCGTCGTCCTTGTAGTCGCAGA (SEQ ID NO:5) and CCTCTGCGACTACAAGGACGACGATGCAAGAAGAAGCGCCAAAGC (SEQ ID NO:6)] were synthesized and used to replace the nucleotide sequences between the StuI and PflMI sites flanking the signal peptidase cleavage site. This reconstituted the C-terminal portion of the signal peptide and newly inserted the 8 amino acid FLAG sequence after this position (FIG. 5). The mature FLAG-tagged MoPrP differs from wild-type MoPrP by the inclusion of the 8 FLAG amino acids, with aspartate, the first amino acid of the FLAG sequence, being amino acid 23 at the N-terminus of the mature FLAG-tagged PrP. A second tagged PrP construct, FLAG-MHM2PrP, was engineered which also inserted the epitope for monoclonal antibody 3F4 derived from SHaPrP (Kascsak et al. (1987) J. Virol. 61:3688–3693) into MoPrP.

The FLAG-MHM2PrP construct was inserted into an MHM2PrP expression cassette previously described by Scott et al. (1992) *Protein Sci.* 1:986–997, cloned into the SPOX.II neo expression vector, and transfected into ScN2A cells (Example 2). Mouse neuroblastoma ScN2A cells which are chronically infected with mouse prions have been previously described (Butler et al. (1988) *J. Virol.* 62:1558–1564).

Neomycin-resistant colonies expressing recombinant PrP were selected in medium containing G418. Mouse PrP is not recognized by monoclonal antibody 3F4, so the inclusion of the 3F4 epitope allows for the discrimination between ectopically expressed PrP and endogenous MoPrP in the event that FLAG-MHM2PrP was not recognized by the anti-FLAG M1 monoclonal antibody.

Western blots of cell lysates from ScN2A cells expressing FLAG-MHM2PrP probed with monoclonal antibody 3F4 showed that inclusion of the FLAG epitope at amino acid 23 does not prevent expression of authentically processed PrP. The FLAG-MHM2PrP appears to be correctly glycosylated, but the apparent molecular weights of FLAG-MHM2PrP glycoforms are about 1–2 kDa greater than MHM2PrP. It is likely that this increase in molecular weight is due to the inclusion of the FLAG peptide in mature PrP^C rather than aberrant processing at the signal peptidase cleavage site since the shifts in size compared to MHM2PrP are in agreement with the predicted molecular weight of the FLAG-PrP fusion protein.

Inclusion of the FLAG sequence into PrP between positions 22 and 23 does not interfere with the processing or biological activity of the tagged protein. The presence of the FLAG epitope did not inhibit proteolytic maturation of PrP at the signal peptidase cleavage site or normal processing and GPI anchorage of PrP at the cell surface. Evidence that the signal peptide is efficiently removed from FLAG-MHM2PrP comes from the immunoreactivity of the ectopically expressed PrP with not only the 3F4 monoclonal antibody but also the anti-FLAG M1 monoclonal antibody, the latter only recognizing fusion proteins with the FLAG tag at the N-terminus. Attempts to detect FLAG-MHM2PrP in transfected ScN2A cells using anti-FLAG M2 monoclonal antibody were unsuccessful. Inclusion of the FLAG tag also does not interfere with proteolytic cleavage at the C-terminus or attachment of PrP on the external surface of cells by GPI anchorage as demonstrated by release of FLAG-MHM2PrP from the cell surface by phosphatidylinositol-specific phospholipase C (PIPLC).

Since placement of the FLAG epitope was required at the N-terminus of mature PrP for recognition by the anti-FLAG M1 monoclonal antibody, it was unknown whether, even though FLAG-MHM2PrP is efficiently expressed and processed, the location of this hydrophilic sequence would interfere with the ability of recombinant PrP to support prion propagation, perhaps by affecting the ability of PrP^C to adopt a conformation essential for the production of infectious prions. A hallmark of PrP^{Sc} is its insolubility in detergents and relative protease resistance.

Proteinase K digestion of PrP in infected ScN2A cells results in the persistence of a core molecule referred to as PrP 27–30 consisting predominantly of amino acid residues 90 to 231. Immunoblotting with anti-FLAG M1 monoclonal antibody failed to detect PrP 27–30 derived from FLAG-MHM2PrP since the FLAG epitope at amino acid 23 is lost following treatment with proteinase K. Using the 3F4 monoclonal antibody to detect proteinase K-resistant FLAG-MHM2PrP, it was found that, like ScN2A cells expressing

MHM2PrP, ScN2A cells expressing FLAG-MHM2PrP efficiently produced PrP 27–30. This phenomenon has also been demonstrated with a nine amino acid peptide sequence, consisting of Ala-Trp-Arg-His-Pro-Gln-Phe-Gly-Gly (SEQ ID NO:3) referred to as the “Strep tag” (Schmidt et al. (1994) *supra*) which behaves in the same way when inserted at the same location. These results establish that the addition of the amino acid sequences at this location of PrP does not interfere with its normal processing, and more importantly, does not interfere with its ability to be converted to the PrP^{Sc} isoform.

The ability of FLAG-MoPrP expressed in transgenic mice to support prion infectivity was tested (Example 4). The FLAG-PrP sequence was engineered into a MoPrP expression cassette which was cloned into the cosSHa.tet cosmid expression vector for transgenic mouse production (Scott et al. (1992) *Protein Sci.* 1:986–997). Five transgenic founders were produced: three in FVB mice and two in FVB/Prn p^{0/0} mice. FVB mice express endogenous MoPrP; FVB/Prn p^{0/0} mice are a line of mice in which the ablated MoPrP gene has been repeatedly backcrossed to FBV and do not express endogenous MoPrP. To simplify analysis, high copy number founders derived only from microinjection of FVB/Prn p^{0/0} embryos were selected for breeding since only transgene-expressed FLAG-MoPrP and not endogenous wild-type MbPrP is expressed in this case. Interference of transgene-directed prion propagation by endogenous wild-type MoPrP has been observed in other experiments (Telling et al. (1994) *Proc. Natl. Acad. Sci.* 91:9936–9940; Telling et al. (1995) *supra*). Using the polyclonal antibody R073, a rabbit polyclonal antibody raised against purified PrP 27–30 of hamster which also reacts with MoPrP, it was estimated by serial dilution and immunodotblotting, that the level of FLAG-MoPrP expression in brain extracts from one line, Tg(FLAG-MoPrP)FVB/Abl 7755, was about 100-fold higher than wild-type levels of MoPrP expression. The extremely high level of FLAG-MoPrP expression meant that FLAG-tagged PrP was more readily detected in brain homogenates of Tg(FLAG-MoPrP)FVB/Abl 7755 mice by the anti-FLAG M1 monoclonal antibody than in ScN2A cells expressing FLAG-MHM2PrP.

In order to determine if Tg(FLAG-MoPrP) mice supported replication of mouse prions, high copy number Tg(FLAG-MoPrP)FVB/Abl 7755 mice were inoculated intracerebrally with mouse RML prions (Example 5). RML is a specific mouse prion isolate derived from Chandler strain mice (Rocky Mountain Laboratory, Hamilton, Mont.). Inoculated mice developed clinical signs of scrapie with an average incubation time of about 52 days (FIG. 6). This is similar to incubation times observed in high copy number Tg(MoPrP)4053 mice, which over-express MoPrP about 8-fold higher than wild-type (Carlson et al. (1994) *Proc. Natl. Acad. Sci.* 91:5690–5694) and is considerably shorter than wild-type non-transgenic mice which have average incubation times of approximately 130 days.

Neuropathological features of the brains of clinically sick include, widespread vacuolation, particularly in the white matter, accompanied by reactive astrocytic gliosis. Hydrolytic autoclaving using polyclonal antibody R073 revealed the presence of PrP-containing plaques predominantly in the corpus callosum.

The brains of clinically sick Tg(FLAG-MoPrP)FVB/Abl 7755 mice inoculated with mouse RML prions contained proteinase K-resistant PrP 27–30 which was detectable using the R073 antibody. Proteinase K treatment results in the loss of the anti-FLAG M1 monoclonal antibody epitope at residue 23 so FLAG-MoPrP^{Sc} is not detected with this

antibody. Since these transgenic mice were derived from FVB/Prn-p^{0/0} mice which express no endogenous MoPrP, the R073-reactive PrP^{Sc} is derived exclusively from transgene-expressed FLAG-MoPrP. The level of FLAG-MoPrP^{Sc} is about 2-fold lower than MoPrP^{Sc} in inoculated wild-type mice. A similar reduction in MoPrP^{Sc} levels is observed in short incubation time Tg(MoPrP)4045 mice which overexpress MoPrP.

Inclusion of the FLAG epitope in MoPrP facilitated the *in situ* detection of PrP^C by histoblot analysis. FLAG-MoPrP^C, but not FLAG-MoPrP^{Sc} was detected using the anti-FLAG M1 monoclonal antibody because the FLAG epitope is lost upon proteinase K digestion of PrP. The distribution of PrP^C was found to be identical using either the anti-FLAG M1 monoclonal antibody or polyclonal R073 antibody. Tg(FLAG-MoPrP) mice inoculated with mouse RML prions revealed a distribution of FLAG-MoPrP^{Sc} which was similar to MoPrP^{Sc} in RML-inoculated wild-type mice.

Since the anti-FLAG M1 monoclonal antibody efficiently recognizes ectopically expressed MoPrP in Tg(FLAG-MoPrP) mice and the levels of FLAG-MoPrP expression were determined to be approximately 100-fold higher than wild-type MoPrP expression, Tg(FLAG-MoPrP)FVB/Abl 7755 is expected to be an excellent source for large amounts of FLAG-MoPrP^C. The presence of the epitope tag allows the tagged PrP molecule to be obtained in high purity under non-denaturing conditions by a one-step immunoaffinity chromatography procedure. Recently, an improved FLAG tag has been generated in which the fifth amino acid of the FLAG sequence was changed from an aspartate residue to a glutamate residue. The binding affinity of the M1 antibody was increased six-fold in Western blots over the original FLAG sequence (Knappik (1994) *Biotechniques* 17:754-761). This raises the possibility that transgenic mice expressing PrP with the improved FLAG tag may be an even better source of recombinant material in the future.

Use of Epitope Placement to Distinguish Between Protein Isoforms

The epitope-tagging system may also be used to differentiate between the conformational shapes of a protein. For example, the epitope tag may be placed in a protein such that the epitope is exposed in one conformational shape and buried in another.

Epitope placement is compared with both epitope-tagged isoforms of MoPrP, HuPrP and MHu2MPrP to locate a placement of the artificial epitope in the protein which results in the exposure of the epitope in the PrP^{Sc} conformational shape and epitope burial in the interior of the PrP^C conformational shape (Example 6). Such tagging approaches are useful in an assay to distinguish between infectious and non-infectious isoforms of PrP. For example, certain regions between residues 109-141 are expected to change from a helix-loop-helix structure into a β -sheet organization. This is expected to change the pattern of presentation of the side chains of these residues so that epitopes that are exposed in the α -helical PrP^C isoform are sequestered in the β -sheet-rich PrP^{Sc} isoform, and vice versa. The exposure of tags placed within this region should be influenced by the conformation change between PrP^C and PrP^{Sc}.

Use of Epitope Tagging to Isolate Replication Intermediates

The epitope tagging technique may be used to isolate "replication intermediates" composed of PrP^C, PrP^{Sc}, and

protein(s) X from ScN2A cells or scrapie-infected mouse brain. FLAG-tagged complexes are immunoprecipitated with anti-FLAG antibodies in the presence of buffer containing low levels of detergent to minimize nonspecific binding and the immunoprecipitates recovered by centrifugation. Proteins from these tagged complexes are released by chelation of the metal ions and analyzed by SDS-PAGE and Western blot. Non-PrP molecules are subjected to N-terminal sequencing leading to the identification of molecular clones encoding such proteins.

This approach of isolating complexes of PrP^C, PrP^{Sc} and protein(s) X has the distinct advantage of being directed at detecting a ternary complex of the three proteins. Complexes of PrP^C or PrP^{Sc} with other proteins may also be detected. This is important for the reconstitution of an *in vivo/in vitro* assays for PrP replication.

EXAMPLES

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use various constructs and perform the various methods of the present invention and are not intended to limit the scope of what the inventors regard as their invention. Unless indicated otherwise, parts are parts by weight, temperature is in degrees centigrade, and pressure is at or near atmospheric pressure. Efforts have been made to ensure accuracy with respect to numbers used, (e.g., length of DNA sequences, molecular weights, amounts, particular components, etc.) but some deviations should be accounted for.

Example 1

Construction of Epitope-Tagged PrP Expression Cassettes

To construct FLAG-tagged PrP expression cassettes, a new recognition sequence for the restriction enzyme *Stu*I was created upstream of the signal peptidase sequence at amino acid 22 (FIG. 4). The *Stu*I site was created by changing a T for an A at nucleotide position 57 in a subclone of the MoPrP gene extending from the start site of the open reading frame to the *Kpn*I site at nucleotide position 277. This was achieved by PCR-mediated mutagenesis in which one of the primers for the reaction contained a mismatch (SEQ ID NO:13): 5'-CCCTCCAGGCTTTGGCCGCTTCTTGACAGAGGCCTACATCAGT-3'.

Complementary 42 and 47 oligonucleotide sequences were synthesized and annealed [(SEQ ID NO:5)(SEQ ID NO:6)]. The nucleotide sequence reconstituted the distal portion of the signal peptide and the signal peptidase cleavage site, followed by the 8 amino acid FLAG sequence and proximal sequence of mature PrP (FIG. 5). This FLAG-tagged sequence was subcloned into two different PrP expression cassettes: MoPrP to produce the FLAG-MoPrP cassette for transgenic mouse production and FLAG-MHM2PrP for expression in cultured cells.

Example 2

FLAG-tagged PrP is Authentically Expressed and is a Substrate for PrP^{Sc} Production in Cultured ScN2A Cells

The FLAG-MHM2PrP expression cassette was cloned into the SPOX.II neo expression vector (Scott et al. (1992) *Protein Sci.* 1:986-997) which allows for direct selection of neomycin resistant transfected cells by growth in medium

containing G418. Mouse neuroblastoma N2A cells chronically infected with mouse prions (ScN2A cells) were grown in 6-well plates and transformed to neomycin resistance by transfection with SPOX-based FLAG-MHM2PrP or control MHM2PrP constructs by DOTAP-mediated transfection (Boehringer Mannheim Biochemicals) and growth in G418-containing medium. Neomycin resistant colonies became apparent approximately 2 weeks after transfection, and these colonies were pooled and passaged.

When stable cultures were established, NP-40 detergent lysates were isolated from either FLAG-MHM2PrP or control MHM2PrP-expressing cells and used for immunoblot analysis. In some experiments, PIPLC was included in the culture medium and PrP cleaved from the cell membrane by the treatment was purified by methanol precipitation and centrifugation.

Analysis of protease-resistant FLAG-MHM2PrP or MHM2PrP in stable cell lines was accomplished by digesting a 1 ml aliquot of cell lysate derived from a confluent 100 mm dish with proteinase K at a concentration of 20 $\mu\text{g}/\text{ml}$ for 1 hour at 37° C. Detergent-resistant, proteinase K-resistant proteins were purified by centrifugation at 40,000 \times g for 1 hour and the pellets resuspended in 20 μl lysis buffer (150 mM NaCl, 10 mM Tris, pH 7.5, 0.5% NP-40, 0.5% deoxycholate). Digested or undigested cell lysates were added to an equal volume of 2 \times SDS-PAGE sample buffer and processed for SDS-PAGE analysis. In all cases, immunoblotting was performed using anti-PrP monoclonal antibody 3F4 directed against hamster PrP^{Sc} (Kascsak et al. (1987) *J. Virol.* 61:3688–3693).

Example 3

Transgenic Mice Expressing FLAG-Tagged PrP

The FLAG-MoPrP ORF expression cassette is flanked by SaII and XhoI, which cleave immediately adjacent to the initiation and termination codons of the PrP ORF, respectively. This allowed for convenient subcloning into the cos.SHaTet expression vector (Scott et al. (1992) *Protein Sci.* 1:986–997) to produce the cos.SHaTet FLAG-MoPrP clone. The isolation and screening of recombinant cosmid clones have been described (Scott et al. (1993) *Cell* 73:979–988, herein specifically incorporated by reference for procedures related to the isolation and screening of recombinant cosmid clones). After verification of the predicted nucleotide sequences of the FLAG-MoPrP ORF, the cosmid NotI fragment, recovered from large scale DNA preparations, was used for microinjections into the pronuclei of fertilized FVB/N or FVB/Prn ^{0/0} mouse embryos as described (Scott et al. (1989) *Cell* 59:847–857; Scott et al. (1992) *supra*, each of which is herein specifically incorporated by reference for procedures for generating transgenic animals).

Five transgenic founders were produced: three in FVB mice and two in FVB/Prn ^{0/0} mice. Genomic DNA, isolated from tail tissue of weanling animals, was screened for the presence of incorporated transgenes using probes that hybridize to the 3'-untranslated region of the SHaPrP gene contained in the cos.SHa.Tet vector (Scott et al. (1992) *supra*). By comparing the hybridization signals of the DNA from weanling mice with standardized DNA samples, it was estimated that one line, Tg(FLAG-MoPrP)FVB/Abl 7755, had transgene copy numbers in excess of approximately 60 transgene copies per cell. Using the polyclonal antibody.

RO73, it was estimated by serial dilution and immunoblotting, that the level of FLAG-MoPrP expression

in brain extracts from the Tg(FLAG-MoPrP)FVB/Abl 7755 line was approximately 100-fold higher than wild-type levels of MoPrP expression.

Example 4

Tg(FLAG-MoPrP) Mice Support Replication of Mouse Prions

The RML isolate from Swiss mice (Chandler (1961) *Lancet* 1:1378–1379) was passaged in Swiss mice from a closed colony at the Rocky Mountain Laboratory (Hamilton, Mont.) or in Swiss CD-1 mice obtained from Charles River Laboratories (Wilmington, Mass.). Mice were inoculated intracerebrally with 30 μl of brain extract using a 27 gauge needle inserted into the right parietal lobe. Beginning 30 days after inoculation, mice were examined for neurologic dysfunction every 3 days. When clinical signs of CNS dysfunction appeared, the mice were examined daily. To confirm the clinical diagnosis, the brains of some animals whose death was obviously imminent were taken for histopathological studies.

Brains were dissected rapidly after sacrifice of the animal and immersion fixed in 10% buffered formalin. The tissue was embedded in paraffin and 8 μm thick histological sections were prepared for staining by the hematoxylin and eosin method and peroxidase immunohistochemical method for glial fibrillary acidic protein and PrP as described previously (DeArmond et al. (1987) *Neurology* 37:1271–1280; Scott et al. (1989) *supra*). Histoblots for localization of PrP^C or protease-resistant PrP were made by pressing 16 mm thick unfixed cryostat sections of brain to nitrocellulose paper as previously described (Taraboulos (1992) *Proc. Natl. Acad. Sci.* 89:7620–7624). To localize PrP^{Sc}, the histoblot was exposed to 400 $\mu\text{g}/\text{ml}$ proteinase K for 18 hours at 37° C. to eliminate PrP^C, exposed to 3 M guanidinium thiocyanate to denature the remaining PrP^{Sc}, followed by immunostaining with PrP specific-antibody RO73 or anti-FLAG M1 monoclonal antibody.

Example 5

Immunopurification of FLAG-MoPrP From Tg (FLAG-MoPrP) Mice

The brain of an uninoculated Tg(FLAG-MoPrP)FVB/Abl 7755 mouse was homogenized in TBS/10 mM CaCl and NP-40 detergent was added to 0.1%. The preparation was applied to a 5 ml column bed comprising anti-FLAG M1 monoclonal antibody coupled to agarose beads. The flow-through was reapplied repeatedly to ensure optimal binding. The column bed was washed three times with 15 ml TBS/Ca. Bound FLAG-MoPrP was eluted with TBS containing 2 mM EDTA which was applied in eight 1 ml aliquots for 10 min each.

Example 6

Use of Epitope Tag to Distinguish Protein Conformational Shapes

The placement of epitope tags at various positions in the prion protein is accomplished by generating recombinant PrP gene sequences harboring the DNA sequence for the epitope at different locations. The working model of PrP^C is the four helix bundle model proposed by Huang et al. (1994) *Proc. Natl. Acad. Sci.* 91:7139–7143. Epitopes are engineered at regions of proposed α -helical structure, which are believed to change conformation during prion replication, or in the loop region connecting the α -helices.

The epitope-tagged sequences are engineered using standard recombinant procedures. The host PrP molecule, MHM2PrP, allows detection by the 3F4 monoclonal antibody (Kascsak et al. (1987) J. Virol. 61:3688–3693) by including the hamster-derived epitope recognized by 3F4. These recombinant PrP constructs are transfected into ScN2A cells which are chronically infected with mouse prions, (Butler et al. (1988) J. Virol. 62:1558–1564). Detection of the ectopically expressed construct is facilitated by 3F4 or by antibodies directed against the heterologous epitope. PrP^{Sc} is detected with 3F4 after digestion with proteinase K. Detection of PrP^{Sc} by 3F4 demonstrates that inclusion of the epitope tag at particular locations does not interfere with PrP^C conversion. Ideally, antibody directed against the epitope tag would detect only PrP^{Sc}, not PrP^C, because in the latter conformation the epitope would be buried and inaccessible to the

antibody. After determination of the ideal location for placement of the epitope tag, transgenic mice expressing PrP with the epitope tag at the desired location are made and infected with prions. Epitope-tagged PrP^{Sc} and PrP^C are isolated from the brains of the transgenic mice using standard procedures. Low resolution analytical techniques such as circular dichroism and infrared spectroscopy are used to determine that the PrP^{Sc} isoform detected by anti-epitope antibody is of the β -sheet conformation, and the PrP^C isoform is of the α -helical conformation.

The instant invention is shown and described herein in what is considered to be the most practical, and preferred embodiments. It is recognized, however, that departures may be made therefrom which are within the scope of the invention, and that modifications will occur to one skilled in the art upon reading this disclosure.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 13

<210> SEQ ID NO 1
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: FLAG peptide

<400> SEQUENCE: 1

Asp Tyr Lys Asp Asp Asp Asp Lys
 1 5

<210> SEQ ID NO 2
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: FLAG peptide

<400> SEQUENCE: 2

Asp Tyr Lys Asp Glu Asp Asp Lys
 1 5

<210> SEQ ID NO 3
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Strep peptide

<400> SEQUENCE: 3

Ala Trp Arg His Pro Gln Phe Gly Gly
 1 5

<210> SEQ ID NO 4
 <211> LENGTH: 6
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: poly-histidine peptide

<400> SEQUENCE: 4

His His His His His His
 1 5

<210> SEQ ID NO 5

-continued

<211> LENGTH: 42
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthesized oligonucleotide

 <400> SEQUENCE: 5

 ttggccgctt cttcttgtca tcgtcgtcct tgtagtcgca ga 42

<210> SEQ ID NO 6
 <211> LENGTH: 47
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthesized oligonucleotide

 <400> SEQUENCE: 6

 cctctgcgac tacaaggacg acgatgacaa gaagaagcgg ccaaagc 47

<210> SEQ ID NO 7
 <211> LENGTH: 254
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

 <400> SEQUENCE: 7

 Met Ala Asn Leu Gly Tyr Trp Leu Leu Ala Leu Phe Val Thr Met Trp
 1 5 10 15
 Thr Asp Val Gly Leu Cys Lys Lys Arg Pro Lys Pro Gly Gly Trp Asn
 20 25 30
 Thr Gly Gly Ser Arg Tyr Pro Gly Gln Gly Ser Pro Gly Gly Asn Arg
 35 40 45
 Tyr Pro Pro Gln Gly Gly Thr Trp Gly Gln Pro His Gly Gly Gly Trp
 50 55 60
 Gly Gln Pro His Gly Gly Ser Trp Gly Gln Pro His Gly Gly Ser Trp
 65 70 75 80
 Gly Gln Pro His Gly Gly Gly Trp Gly Gln Gly Gly Gly Thr His Asn
 85 90 95
 Gln Trp Asn Lys Pro Ser Lys Pro Lys Thr Asn Leu Lys His Val Ala
 100 105 110
 Gly Ala Ala Ala Ala Gly Ala Val Val Gly Gly Leu Gly Gly Tyr Met
 115 120 125
 Leu Gly Ser Ala Met Ser Arg Pro Met Ile His Phe Gly Asn Asp Trp
 130 135 140
 Glu Asp Arg Tyr Tyr Arg Glu Asn Met Tyr Arg Tyr Pro Asn Gln Val
 145 150 155 160
 Tyr Tyr Arg Pro Val Asp Gln Tyr Ser Asn Gln Asn Asn Phe Val His
 165 170 175
 Asp Cys Val Asn Ile Thr Ile Lys Gln His Thr Val Thr Thr Thr Thr
 180 185 190
 Lys Gly Glu Asn Phe Thr Glu Thr Asp Val Lys Met Met Glu Arg Val
 195 200 205
 Val Glu Gln Met Cys Val Thr Gln Tyr Gln Lys Glu Ser Gln Ala Tyr
 210 215 220
 Tyr Asp Gly Arg Arg Ser Ser Ser Thr Val Leu Phe Ser Ser Pro Pro
 225 230 235 240
 Val Ile Leu Leu Ile Ser Phe Leu Ile Phe Leu Ile Val Gly
 245 250

-continued

```

<210> SEQ ID NO 8
<211> LENGTH: 253
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8
Met Ala Asn Leu Gly Cys Trp Met Leu Val Leu Phe Val Ala Thr Trp
 1          5          10          15
Ser Asp Leu Gly Leu Cys Lys Lys Arg Pro Lys Pro Gly Gly Trp Asn
 20          25          30
Thr Gly Gly Ser Arg Tyr Pro Gly Gln Gly Ser Pro Gly Gly Asn Arg
 35          40          45
Tyr Pro Pro Gln Gly Gly Gly Gly Trp Gly Gln Pro His Gly Gly Gly
 50          55          60
Trp Gly Gln Pro His Gly Gly Gly Trp Gly Gln Pro His Gly Gly Gly
 65          70          75          80
Trp Gly Gln Pro His Gly Gly Gly Trp Gly Gln Gly Gly Gly Thr His
 85          90          95
Ser Gln Trp Asn Lys Pro Ser Lys Pro Lys Thr Asn Met Lys His Met
 100         105         110
Ala Gly Ala Ala Ala Gly Ala Val Val Gly Gly Leu Gly Gly Tyr
 115         120         125
Met Leu Gly Ser Ala Met Ser Arg Pro Ile Ile His Phe Gly Ser Asp
 130         135         140
Tyr Glu Asp Arg Tyr Tyr Arg Glu Asn Met His Arg Tyr Pro Asn Gln
 145         150         155
Val Tyr Tyr Arg Pro Met Asp Glu Tyr Ser Asn Gln Asn Asn Phe Val
 165         170         175
His Asp Cys Val Asn Ile Thr Ile Lys Gln His Thr Val Thr Thr Thr
 180         185         190
Thr Lys Gly Glu Asn Phe Thr Glu Thr Asp Val Lys Met Met Glu Arg
 195         200         205
Val Val Glu Gln Met Cys Ile Thr Gln Tyr Glu Arg Glu Ser Gln Ala
 210         215         220
Tyr Tyr Gln Arg Gly Ser Ser Met Val Leu Phe Ser Ser Pro Pro Val
 225         230         235         240
Ile Leu Leu Ile Ser Phe Leu Ile Phe Leu Ile Val Gly
 245         250

```

```

<210> SEQ ID NO 9
<211> LENGTH: 263
<212> TYPE: PRT
<213> ORGANISM: bovine sp.

```

```

<400> SEQUENCE: 9
Met Val Lys Ser His Ile Gly Ser Trp Ile Leu Val Leu Phe Val Ala
 1          5          10          15
Met Trp Ser Asp Val Gly Leu Cys Lys Lys Arg Pro Lys Pro Gly Gly
 20          25          30
Trp Asn Thr Gly Gly Ser Arg Tyr Pro Gly Gln Gly Ser Pro Gly Gly
 35          40          45
Asn Arg Tyr Pro Pro Gln Gly Gly Gly Gly Trp Gly Gln Pro His Gly
 50          55          60
Gly Gly Trp Gly Gln Pro His Gly Gly Gly Trp Gly Gln Pro His Gly
 65          70          75          80
Gly Gly Trp Gly Gln Pro His Gly Gly Gly Trp Gly Gln Pro His Gly

```

-continued

85				90				95							
Gly	Gly	Gly	Trp	Gly	Gln	Gly	Gly	Thr	His	Gly	Gln	Trp	Asn	Lys	Pro
			100												110
Ser	Lys	Pro	Lys	Thr	Asn	Met	Lys	His	Val	Ala	Gly	Ala	Ala	Ala	Ala
			115				120								125
Gly	Ala	Val	Val	Gly	Gly	Leu	Gly	Gly	Tyr	Met	Leu	Gly	Ser	Ala	Met
			130				135								140
Ser	Arg	Pro	Leu	Ile	His	Phe	Gly	Ser	Asp	Tyr	Glu	Asp	Arg	Tyr	Tyr
							150								160
Arg	Glu	Asn	Met	His	Arg	Tyr	Pro	Asn	Gln	Val	Tyr	Tyr	Arg	Pro	Val
															175
Asp	Gln	Tyr	Ser	Asn	Gln	Asn	Asn	Phe	Val	His	Asp	Cys	Val	Asn	Ile
			180												190
Thr	Val	Lys	Glu	His	Thr	Val	Thr	Thr	Thr	Thr	Lys	Gly	Glu	Asn	Phe
			195												205
Thr	Glu	Thr	Asp	Ile	Lys	Met	Met	Glu	Arg	Val	Val	Glu	Gln	Met	Cys
			210				215								220
Val	Thr	Gln	Tyr	Gln	Lys	Glu	Ser	Gln	Ala	Tyr	Tyr	Asp	Gln	Gly	Ala
															240
Ser	Val	Ile	Leu	Phe	Ser	Ser	Pro	Pro	Val	Ile	Leu	Leu	Ile	Ser	Phe
															255
Leu	Ile	Phe	Leu	Ile	Val	Gly									
			260												

<210> SEQ ID NO 10

<211> LENGTH: 255

<212> TYPE: PRT

<213> ORGANISM: ovine sp.

<400> SEQUENCE: 10

Met	Val	Lys	Ser	His	Ile	Gly	Ser	Trp	Ile	Leu	Val	Leu	Phe	Val	Ala
															15
Met	Trp	Ser	Asp	Val	Gly	Leu	Cys	Lys	Arg	Pro	Lys	Pro	Gly	Gly	
			20												30
Trp	Asn	Thr	Gly	Gly	Ser	Arg	Tyr	Pro	Gly	Gln	Gly	Ser	Pro	Gly	Gly
			35				40								45
Asn	Arg	Tyr	Pro	Pro	Gln	Gly	Gly	Gly	Trp	Gly	Gln	Pro	His	Gly	
			50				55								60
Gly	Gly	Trp	Gly	Gln	Pro	His	Gly	Gly	Gly	Trp	Gly	Gln	Pro	His	Gly
			65				70								80
Gly	Ser	Trp	Gly	Gln	Pro	His	Gly	Gly	Gly	Trp	Gly	Gln	Gly	Gly	
			85												95
Ser	His	Ser	Gln	Trp	Asn	Lys	Pro	Ser	Lys	Pro	Lys	Thr	Asn	Met	Lys
			100												110
His	Val	Ala	Gly	Ala	Ala	Ala	Ala	Gly	Ala	Val	Val	Gly	Gly	Leu	Gly
			115				120								125
Gly	Tyr	Met	Leu	Gly	Ser	Ala	Met	Ser	Arg	Pro	Leu	Ile	His	Phe	Gly
			130				135								140
Asn	Asp	Tyr	Glu	Asp	Arg	Tyr	Tyr	Arg	Glu	Asn	Met	Tyr	Arg	Tyr	Pro
			145				150								160
Asn	Gln	Val	Tyr	Tyr	Arg	Pro	Val	Asp	Gln	Tyr	Ser	Asn	Gln	Asn	Asn
			165												175
Phe	Val	His	Asp	Cys	Val	Asn	Ile	Thr	Val	Lys	Gln	His	Thr	Val	Thr
			180												190

-continued

```

Thr Thr Thr Lys Gly Glu Asn Phe Thr Glu Thr Asp Ile Lys Ile Met
    195                200                205

Glu Arg Val Val Glu Gln Met Cys Ile Thr Gln Tyr Gln Arg Glu Ser
    210                215                220

Gln Ala Tyr Tyr Gln Arg Gly Ala Ser Val Ile Leu Phe Ser Ser Pro
    225                230                235                240

Pro Val Ile Leu Leu Ile Ser Phe Leu Ile Phe Leu Ile Val Gly
    245                250                255

```

<210> SEQ ID NO 11
 <211> LENGTH: 13
 <212> TYPE: PRT
 <213> ORGANISM: Influenza virus

<400> SEQUENCE: 11

```

Tyr Pro Tyr Asp Val Pro Asp Tyr Ala Ile Glu Gly Arg
  1          5          10

```

<210> SEQ ID NO 12
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

```

Glu Gln Lys Leu Leu Ser Glu Glu Asp Leu Asn
  1          5          10

```

<210> SEQ ID NO 13
 <211> LENGTH: 42
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthesized oligonucleotide

<400> SEQUENCE: 13

ccctccaggc ttggccgct tcttgacagag gctacatca gt

42

What is claimed is:

1. A recombinant nucleic acid construct comprising:
 - a first nucleic acid sequence encoding a PrP protein; and
 - a second nucleic acid sequence encoding a heterologous epitope domain;
 wherein the PrP protein has two different three dimensional conformations, a first PrP^C and a second PrP^{Sc} conformation, wherein the heterologous epitope domain is spatially positioned within the PrP protein such that the epitope domain is at least 20% more exposed in the second conformation relative to the first conformation.
2. The construct of claim 1, wherein the PrP protein is a human PrP protein.
3. The construct of claim 1, wherein the PrP protein is selected from the group consisting of a natural, synthetic, and chimeric PrP protein.
4. The construct of claim 1, wherein the heterologous epitope domain is a peptide selected from the peptide group of FLAG, Strep, poly-histidine, human c-myc peptide recognized by monoclonal antibody 9E10, hemagglutinin peptide recognized by monoclonal antibody 12CA5, and Glu-Glu-Phe.
5. The construct of claim 4, wherein said FLAG peptide has a sequence Asp-Tyr-Lys-Asp-Asp-Asp-Asp-Lys (SEQ ID NO:1) or Asp-Tyr-Lys-Asp-Glu-Asp-Asp-Lys (SEQ ID NO:2).
6. The construct of claim 4, wherein said Strep peptide has the sequence Ala-Trp-Arg-His-Pro-Gln-Phe-Gly-Gly (SEQ ID NO:3).
7. The construct of claim 4, wherein said poly-histidine peptide has the sequence His-HisHis-His-His-His (SEQ ID NO:4).
8. The construct of claim 4, wherein said c-myc peptide has the sequence Glu-Gln-LysLeu Leu-Ser-Glu-Glu-Asp-Leu-Asn (SEQ ID NO:12).
9. The construct of claim 4, wherein said hemagglutinin peptide has the sequence Tyr-ProTyr-Asp-Val-Pro-Asp-Tyr-Ala-Ile-Glu-Gly-Arg (SEQ ID NO: 11).
10. The construct of claim 1, wherein the PrP protein has a heterologous FLAG epitope domain inserted before amino acid residue 23 of the PrP sequence, wherein said amino acid residue of the PrP protein corresponds to amino acid residue 23 of mouse PRP (SEQ ID NO: 7).
11. A cell line having operatively inserted therein a recombinant nucleic acid construct comprising:
 - a first nucleic acid sequence encoding a PrP protein; and
 - a second nucleic acid sequence encoding a heterologous epitope domain;
 wherein the PrP protein has two different three dimensional conformations, a first PrP^C and a second PrP^{Sc} conformation, wherein the heterologous epitope domain is spatially positioned within the PrP protein

31

such that the epitope domain is at least 20% more exposed in the second conformation relative to the first conformation.

12. An assay for detecting infections prior material, said assay comprising the steps of:

a) generating a cell line expressing an epitome tagged PrP protein, said cell line having an operatively inserted therein a recombinant nucleic acid construct comprising:

a first nucleic acid sequence encoding a PrP protein;

and

a second nucleic acid sequence encoding a heterologous epitope domain;

wherein the PrP protein has two different three dimensional conformations, a first PrP^C and a second PrP^{Sc} conformation, wherein the heterologous epitope domain is spatially positioned within the PrP protein such that the epitope domain is at least 20% more exposed in the second conformation relative to the first conformation;

b) inoculating said cell with material containing infectious prion protein; and

32

c) detecting the presence of epitope-tagged PrP^{Sc} with an antibody that binds to the heterologous epitope.

13. The assay of claim 12, wherein said PrP protein is a cow protein, and the inoculating material is from a cow.

14. The assay of claim 12, wherein said PrP protein is a chimer cow/mouse protein, and the inoculating material is from a cow.

15. The assay of claim 12, wherein said PrP protein is a human protein, and the inoculating material is from a human.

16. The assay of claim 12, wherein said PrP protein is a chimeric human/mouse protein, and the inoculating material is from a cow.

17. The assay of claim 12, wherein said PrP protein is a sheep protein, and the inoculating material is from a cow.

18. The assay of claim 12, wherein said PrP protein is a chimeric sheep/mouse protein, and the inoculating material is from a cow.

* * * * *